

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/000514

International filing date: 20 January 2005 (20.01.2005)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 04001175.1
Filing date: 21 January 2004 (21.01.2004)

Date of receipt at the International Bureau: 23 March 2005 (23.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04001175.1

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 04001175.1
Demande no:

Anmeldetag:
Date of filing: 21.01.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Newron Pharmaceuticals S.p.A.
Via Ariosto, 21
20091 Bresso (MI)
ITALIE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Alfa-aminoamide derivatives useful in the treatment of lower urinary tract disorders

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K31/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

BIANCHETTI • BRACCO • MINOJA
STUDIO CONSULENZA BREVETTUALE

**PLEASE ACKNOWLEDGE SAFE RECEIPT OF
THIS APPLICATION BY FAX**


The fee for acknowledgement of receipt
will be paid in due time

Thank You!

04 001 175 . 1

OUR REF: SCB 1337 EUR

FAX Received on 21/01/04 with 45
pages & has been given the above no.





From: MINOJA, Fabrizio, Dr.
BIANCHETTI BRACCO MINOJA SRL
Via Rossini, 8
20122 MILANO MI (Italy)

Telefax n. (02) 783078
Tel (02) 76021218 - 76021192

To: EUROPEAN PATENT OFFICE
Erhardtstraße 27
D-80298 MÜNCHEN
Germany

This transmission consists of ...**45**..... pages, including this cover page.

If the transmission is not clear, please call or fax.



Request for grant of a European Patent:

**ALFA-AMINOAMIDE DERIVATIVES USEFUL IN THE
TREATMENT OF LOWER URINARY TRACT DISORDERS**

in the name of: **NEWRON PHARMACEUTICALS S.p.A.**

Our ref.: **SCB 1337 EUR/FM/cgr**

The confirmation copy follows by DHL

Milano, 21 January 2004

ALFA-AMINOAMIDE DERIVATIVES USEFUL IN THE TREATMENT OF LOWER URINARY TRACT DISORDERS

FIELD OF THE INVENTION

The invention relates to novel and known α -aminoamide derivatives, a chemical class of sodium channel blockers, and their use for treating lower urinary tract disorders and to pharmaceutical compositions containing them.

BACKGROUND OF THE INVENTION

Lower urinary tract (LUT) disorders, that affect the quality of life of millions of people in the world every year, include but are not limited to overactive bladder (OAB), prostatitis and prostatic dysplasia, interstitial cystitis, benign prostatic hyperplasia and urinary incontinence.

OAB is the most common term currently used in clinical medicine to describe a complex of lower urinary tract symptoms with or without incontinence. The symptoms usually include urgency, frequency, nocturia, troublesome or incomplete emptying and occasionally pain. Causes of bladder overactivity include neurological illness or injury, bladder outlet obstruction, urethral weakness, detrusor hyperactivity and impaired contractility in elderly patients, emergence of new voiding reflexes and so-called idiopathic bladder overactivity. Neurogenic overactive bladder occurs as the results of neurological damage due to disorders such as stroke, Parkinson's disease, diabetes, multiple sclerosis, peripheral neuropathy or spinal cord lesions. By contrast, non-neurogenic OAB can result from non-neurological abnormalities including bladder stones, muscle disease, urinary tract infection or drug side effects. OAB may result from hypersensitivity of sensory neurons of the urinary bladder, arising from various factors including inflammatory conditions, hormonal imbalances, and prostate hypertrophy.

Prostatitis is a term to describe inflammatory conditions of the prostate.

gland. It is thought that most cases of prostatitis result from bacterial infection, but evidence of infection is not always found. An infected or inflamed prostate can cause painful urination with serious complications. There are four types of prostatitis: acute bacterial prostatitis, chronic bacterial
5 prostatitis, non-bacterial prostatitis and prostatic dyspareunia, also called chronic pelvic pain syndrome, that is a condition associated with painful symptoms of chronic non-bacterial prostatitis without an inflammation of the prostate.

Interstitial cystitis (IC) is a chronic inflammatory condition of the bladder that causes frequent, urgent and painful urination and pelvic
10 discomfort. Unlike common cystitis (inflammation of the bladder caused by bacterial infection), no infectious agent has been found in IC.

Benign prostatic hyperplasia (BPH) is a non-malignant enlargement of the prostate that is very common in men over 40 years. The prostate grows in two different ways. In one type of growth, cells multiply around the urethra, in
15 the second cells grow into the urethra and the bladder outlet area. This second type of growth typically requires surgery. Common symptoms of BPH include: blood in the urine, feeling that bladder has not emptied completely after the urination, frequent urination, nocturia, urgent need to urinate, incontinence. In severe cases of BPH another symptom is the acute urinary
20 retention.

The function of the LUT is to store and periodically release urine. This requires the orchestration of storage and micturition reflexes involving both the sympathetic and parasympathetic components of the autonomic nervous system and motor pathways.

25 Urinary incontinence (UI) occurs when the lower urinary tract does not store urine properly and there is involuntary loss of urine. There are 3 types of UI: urge, stress and mixed. Urge UI is considered to be due to an overactive bladder, while stress UI is considered to be due to decreased urethral outlet

resistance. Mixed UI contains both components. Symptoms are: urinary frequency, urgency, nocturia, accidental loss of urine. When conscious control of the parasympathetic micturition reflex is compromised, conditions of OAB and/or UI arise, creating serious health and social concerns.

- 5 Drugs that suppress the micturition reflex could be useful for treating OAB and UI. One of the target for suppressing the OAB consists in the primary afferent neurons and their peripheral terminals in the bladder. Therapeutic agents that suppress action potential initiation and/or propagation along primary afferent fibers, through modulation of ion channels, would be
10 expected to increase the volume threshold for activation of the micturition reflex and thus reduce bladder overactivity and urgency.

Current treatments for OAB include antimuscarinic drugs, diet modification, programs in bladder training, electrical stimulation and surgery.

- 15 Currently, there are no established treatments for prostatitis and prostatic dysphasia. Antibiotics, COX-2 inhibitors and α -adrenergic blockers are often prescribed, but their efficacy has not been established.

Treatments for IC that include the administration of antihistamines, sodium pentosanpolysulfate, dimethylsulfoxide, steroids, tricyclic antidepressants and narcotic antagonists have generally been unsuccessful.

- 20 Non-invasive treatments for BPH include androgen deprivation therapy and the use of 5 α -reductase inhibitors and α -adrenergic blockers with minimal efficacy.

- 25 Because existing therapies and treatments for LUT disorders are associated with limitations as described above, new therapies and treatments are desirable.

However, despite the large number of available therapeutic agents, their use suffers of limited efficacy and side effects such as dry mouth, dry eyes, drowsiness, palpitations, cardiac rhythm alteration, drowsiness, mental

confusion particularly in the elderly and constipation which have proven difficult for individuals to tolerate.

Since sensory afferent neurons, expressing sodium channels, are hypersensitized in bladder disfunctions, it seems reasonable to assume that blocking sodium channels activity may represent a novel strategy for the treatment of lower urinary tract disorders.


The present invention provides rapid and highly effective methods for treating a variety of lower urinary tract disorders by utilizing, *in vivo*, certain α -aminoamide compounds of the invention in a therapy as an alternative to existing treatments.

WO90/14334, WO94/22808, WO97/05102, WO97/0511 and WO99/35125. The text of which are herein incorporated by reference, disclose substituted benzylaminopropionamide compounds active on the central nervous system and useful as anti-epileptic, anti-Parkinson, neuroprotective, antidepressant and antispastic hypnotic agents (*see also* Pevarello P. *et al.* (1998), "Synthesis and anticonvulsant activity of a new class of 2-[(arylalkyl)amino]alkanamide derivatives", *J. Med. Chemistry*, 41: 579-590). WO99/35125 and WO99/35123 disclose substituted benzylaminopropanamide compounds active on the central nervous system and useful as analgesic agents (*see also* Veneroni O. *et al.* (2003) "Anti-allodynic effect of NW-1029, a novel Na⁺ channel blocker, in experimental animal models of inflammatory and neuropathic pain", *Pain* 102(1-2):17-25).

DETAILED DESCRIPTION OF THE INVENTION

The invention concerns the use of compounds of Formula I for the preparation of a medicament for treating lower urinary tract disorders. The compounds according to the invention are α -aminoamide derivatives of formula (I):

5



(I)

wherein:

- R is a furyl, thienyl, or pyridyl ring or a phenyl ring, optionally substituted by one or two substituents independently selected from halogen, hydroxy, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy or trifluoromethyl;
- R₁ is hydrogen or C₁-C₆ alkyl or C₃-C₇ cycloalkyl;
- R₂ and R₃ are independently selected from hydrogen; C₁-C₄ alkyl, optionally substituted by hydroxy or phenyl, phenyl, the phenyl rings being optionally substituted by one or two substituents independently selected from C₁-C₆ alkyl, halogen, hydroxy, C₁-C₆ alkoxy or trifluoromethyl; or R₂ and R₃, taken with the carbon atom which they are linked to, form a C₃-C₆ cycloalkyl ring;
- R₄, R₅ are, independently, hydrogen, C₁-C₆ alkyl or C₃-C₇ cycloalkyl; or R₄ and R₅, taken together with the nitrogen atom they are linked to, form a 5-7 atom saturated heterocyclic ring;
- X is CH₂, O, S;
- Y, Z are hydrogen or taken together form a 5-7 saturated or unsaturated carbocycle or a heterocycle

or isomers, mixtures, and pharmaceutically acceptable salts thereof.

The alkyl and alkoxy groups can be branched or straight.

Pharmaceutically acceptable salts of the compounds of the invention include, for example, acid addition salts with inorganic acids, e.g., nitric, hydrochloric, hydrobromic, sulfuric and phosphoric acids and the like, or

organic acids, e.g., acetic, propionic, glycolic, lactic, oxalic, malonic, malic, tartaric, citric, succinic, benzoic, cinnamic, mandelic, methanesulfonic, p-toluenesulfonic and salicylic acids, and the like.

Some of the compounds of formula (I) can have asymmetric carbon atoms, and therefore can exist either as racemic mixtures or as individual optical isomers (enantiomers). Accordingly, the term "pharmaceutically acceptable salts" of the α -aminoamide of formula (I) is also meant to include within its scope all the possible isomers and their mixtures, and any pharmaceutically acceptable metabolite, bioprecursor and/or pro-drug, i.e., a compound which has a structural formula different from the one of the the α -aminoamide of formula (I), and yet is directly or indirectly converted *in vivo* into a compound having formula (I), upon administration to a mammal, particularly a human being.

X and Y are preferably in reciprocal meta or para position.

The compounds of formula (I) wherein Y together with Z form a dihydrobenzofuran or dihydrobenzothiophen or a dihydrobenzo(thio)pyran or a tetrahydrobenz(thio)oxepin heterocycle are new and are a further object of the invention.

Preferably, in the compounds of formula I, R is a phenyl ring, optionally substituted by one or two substituents independently selected from a halogen, trifluoromethyl, a methoxy, or a thienyl ring;

- R₁ is hydrogen or C₁-C₄ alkyl;
- One of R₂ and R₃ is hydrogen and the other is C₁-C₄ alkyl, optionally substituted by hydroxy or phenyl, optionally substituted by one or two halogen atoms, or R₂ and R₃ are both methyl, or together they form with the atom they are linked to a cyclopropyl or a cyclopentyl ring; and
- R₄, R₅ are hydrogen or C₁-C₄ alkyl, or together with the nitrogen

they are linked to, form a pyrrolidine or piperidine ring, and the pharmaceutically acceptable salts thereof.

Examples of compounds for use according to the invention include:

- 2-(4-Phenethyl-benzylamino)-propanamide;
- 5 2-(4-Benzyloxybenzylamino)-propanamide;
- 2-(3-Benzyloxybenzylamino)-propanamide;
- 2-(4-Benzylthiobenzylamino)-propanamide;
- 2-(4-Benzyloxy-benzylamino)-3-N-dimethyl-butanamide;
- 2-[4-(2-Methoxybenzyloxy)-benzylamino]-propanamide;
- 10 2-[4-(2-Fluorobenzyloxy)-benzylamino]-propanamide;
- 2-[3-(2-Fluorobenzyloxy)-benzylamino]-propanamide;
- 2-[4-(2-Fluorobenzyloxy)-benzylamino]-propanamide;
- 2-[4-(2-Fluorobenzyloxy)-benzylamino]-2-methyl-propanamide;
- 2-[4-(2-Fluorobenzyloxy)-benzylamino]-N-methyl-propanamide;
- 15 2-[3-(3-Fluorobenzyloxy)-benzylamino]-propanamide;
- 2-[4-(3-Fluorobenzyloxy)-benzylamino]-propanamide;
- 2-[4-(3-Methoxybenzyloxy)-benzylamino]-propanamide;
- 2-[4-(3-Cyanobenzyloxy)-benzylamino]-propanamide;
- 2-[4-(3-Fluorobenzyloxy)-benzylamino]-propanamide;
- 20 2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-methyl-propanamide;
- 2-[4-(3-Fluorobenzyloxy)-benzylamino]-N-methyl-propanamide;
- 2-[4-(4-Fluorobenzyloxy)-benzylamino]-propanamide;
- 2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-methyl-propanamide;
- 2-[4-(2-Chlorobenzyloxy)-benzylamino]-propanamide;
- 25 2-[4-(3-Chlorobenzyloxy)-benzylamino]-propanamide;
- 2-(4-Benzyloxybenzylamino)-3-hydroxy-propanamide;
- 2-[4-(3-Fluorobenzyloxy)-benzylamino]-3-hydroxy-propanamide;
- 2-[4-(3-Fluorobenzyloxy)-benzylamino]-3-hydroxy-propanamide;

- 2-(4-Benzyloxybenzylamino)-3-hydroxy-N-methyl-propanamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-3-hydroxy-N-methyl-
propanamide;
2-[4-(3-Fluorobenzyloxy)-benzylamino]-3-hydroxy-N-methyl-
5 propanamide;
2-[4-(2-Chlorobenzyloxy)-benzylamino]-3-hydroxy-N-methyl-
propanamide;
2-[4-(3-Cyanobenzyloxy)-benzylamino]-3-hydroxy-N-methyl-
propanamide;
10 2-[4-(3-Cyanobenzyloxy)-benzylamino]-2-methyl-3-hydroxy-N-methyl-
propanamide;
2-[4-(3-Chlorobenzyloxy)-phenylethylamino]-propanamide;
2-[4-[2-(3-Fluorophenyl)-ethyloxy]benzylamino]-propanamide;
2-[4-[2-(3-Fluorophenyl)-ethyl]benzylamino]-propanamide;
15 2-[N-(4-Benzyloxybenzyl)-N-methylamino]-propanamide;
2-[4-[(3-Chlorobenzyloxy)-phenylethyl]-amino]-propanamide;
2-[4-Benzylthiobenzylamino]-propanamide;
2-[4-(2-Fluorobenzylthio)-benzylamino]-propanamide;
2-[4-(3-Fluorobenzylthio)-benzylamino]-propanamide;
20 2-[4-(3-Phenylpropyloxy)-benzylamino]-propanamide;
2-[4-(4-Phenylbutyloxy)-benzylamino]-propanamide;
2-[4-(5-Phenylpentyloxy)-benzylamino]-propanamide;
2-(4-Benzyloxybenzylamino)-3-phenyl-N-methyl-propanamide;
2-(4-Benzyloxybenzylamino)-3-methyl-N-methyl-butanamide;
25 2-(4-Benzyloxybenzylamino)-2-phenyl-acetamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-2-phenyl-acetamide;
2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-phenyl-acetamide;
2-[4-(2-Fluorobenzyloxy)-benzyl-N-methylamino]-2-phenyl-acetamide;

2-[4-(3-Fluorobenzyloxy)-benzyl-N-methylamino]-2-phenyl-acetamide;

2-[4-(3-Chlorobenzyloxy)-benzylamino]-2-phenyl-acetamide;

2-[4-(2-Fluorobenzyloxy)-benzylamino]-2-(2-fluorophenyl)-acetamide;

2-[4-(2-Fluorobenzyloxy)-benzylamino]-2-(3-fluorophenyl)-acetamide;

5 2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-(2-fluorophenyl)-acetamide;

2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-(3-fluorophenyl)-acetamide;

2-[4-(3-Chlorobenzyloxy)-benzylamino]-2-(3-fluorophenyl)-acetamide;

2-(4-(2-Thienyloxy)-benzylamino)-propanamide;

Examples of new compounds of formula (I) include:

10 2-[(3-Benzyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-
propanamide;

2-[(3-Benzyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-
propanamide;

2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
15 amino)-propanamide;

2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-N-methyl-propanamide;

2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-propanamide;

20 2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-N-methyl-propanamide;

2-[(3-Phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-
propanamide;

2-[(3-Phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-
25 propanamide;

2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-propanamide;

2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-

amino)-N-methyl-propanamide;

2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-propanamide;

2-{3-[2-(3-Chloro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
5 amino)-propanamide;

2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-N-methyl-propanamide;

2-[(3-Phenethyl-2,3-dihydro-benzopyran-6-ylmethyl)-amino]-
propanamide;

10 2-[(4-Phenethyl-2,3-dihydro-benzoxepin-7-ylmethyl)-amino]-
propanamide;

2-[(3-Benzyl-2,3-dihydro-benzothiophen-5-ylmethyl)-amino]-
propanamide;

2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
15 amino)-propanamide;

2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide;

2-[(3-Phenethyl-2,3-dihydro-benzothiophen-5-ylmethyl)-amino]-
propanamide;

20 2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide;

2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide;

25 2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-N-methyl-propanamide

or isomers, mixtures, and pharmaceutically acceptable salts thereof.

Preferred compound of formula (I), which can be used singly, or in
combination with other compounds of formula (I) are (S)-(+)-2-[3-(2-

- fluorobenzyloxy)-benzylamino]-propanamide; (S)-(+)-2-[4-(2-fluorobenzyloxy)-benzylamino]-propanamide (NW-1029), (S)-(+)-2-[4-(2-fluorobenzyloxy)-benzylamino]-N-methyl-propanamide (S)-(+)-2-[3-(3-fluorobenzyloxy)-benzylamino]-propanamide, (S)-(+)-2-[4-(3-fluorobenzyloxy)-benzylamino]-propanamide, (S)-(+)-2-[4-(3-fluorobenzyloxy)-benzylamino]-N-methyl-propanamide, (S)-(+)-2-(4-phenethyl-benzylamino)-propanamide, (R)-(-)-2-(4-benzyloxy-benzylamino)-3-N-dimethyl-butanamide, (S)-(+)-2-(4-benzylthiobenzylamino)-propanamide, and 2-[(3-phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-propanamide and (2R/3'S,R)-2-[(3-phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-propanamide, 2-{3-[2-(2-fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-propanamide, 2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-propanamide, (2R/3'S,R)-2-{3-[2-(2-fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-propanamide, (2R/3'S,R)-2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-propanamide.

The preparation of the known compounds of formula I is disclosed in WO90/14334, WO94/22808, WO97/05102, WO97/0511, WO99/35125 and WO99/35123.

- The preparation of the new compounds may be carried out by reacting an ester of 4-hydroxy-3-iodo-benzoic acid with the suitable substituted 4-phenylbut-2-enyl bromides or ethyl cinnamyl bromides in the presence of bases (e.g. NaH) and a suitable catalyst such as a crown ether. The obtained 3-iodo-4-[(2E)-4-phenyl-but-2-enyl]-oxy}benzoate or 3-benzyl-2,3-dihydro-1-benzofuran-5-carboxylate esters are then transformed into the corresponding 3-(2-phenylethyl)-2,3-dihydro-1-benzofuran-5-carboxylates or 3-benzyl-2,3-dihydro-1-benzofuran-5-carboxylates by ring closure in the presence of AIBN and triethylamine. The reduction by LiAlH₄ of the 5-carboxylates.

groups to 5-hydroxymethyl groups followed by oxidation to 5-carboxyaldehyde groups provides intermediate compounds which, by reductive amination with an amine of formula $\text{NHR}_1\text{-CR}_2\text{R}_3\text{CONR}_4\text{R}_5$, where $\text{R}_1\text{-R}_5$ have the same meanings reported above, yield the desired benzofuran derivatives. Similarly, starting from 4-mercapto-3-iodo-benzoic acid esters, the corresponding benzothiophen derivatives may be obtained.

In one embodiment the patient being treated is a mammal, including humans, in need of alleviation, prevention, or inhibition of symptoms of the lower urinary tract disorders.

Particularly, the mammal in need of the above mentioned treatment is administered a dose of an α -aminoamide of formula (I) as above defined which ranges from about 0.3 to about 100 mg/kg of body weight per day. "Treatment" as used herein includes any care by procedures or applications to a mammal, and particularly a human, that are intended to a) prevent the disease or disorder from occurring in a subject that may be predisposed to the disease/disorder, but has not yet been diagnosed with having it; b) inhibiting the disease/disorder, or condition, i.e., arresting its development; or c) relieving the disease/disorder, or condition, i.e., causing regression of the disease/disorder, or condition.

Lower urinary tract disorders in a mammal, including humans, can thus be inhibited, alleviated and prevented. Examples of lower urinary tract disorders in mammals which can be treated by administering one or more α -aminoamide compounds of formula (I) include, but are not limited to: overactive bladder (OAB), prostatitis and prostatic dysplasia, interstitial cystitis, benign prostatic hyperplasia, and urinary incontinence. In another aspect, the invention includes an α -aminoamide of formula (I), administered as the active agent of a pharmaceutically acceptable composition having activity in the lower urinary tract disorders, which can be prepared by conventional

procedures known in the art, for instance by mixing the active agent with a pharmaceutically acceptable, therapeutically inert organic and/or inorganic carrier or excipient materials.

"Combination therapy" (or "co-therapy") includes the administration of an alpha-aminoamide compound of formula (I) of the invention and at least a second agent as part of a specific treatment regimen intended to provide the beneficial effect from the co-action of these therapeutic agents. Benefits of such combinations include reduction of the dose of conventional therapeutic agents (*i.e.*, other than the agents of the present invention) with consequent reduction of the side-effects of such conventional agents. The beneficial effect of the combination includes, but is not limited to, pharmacokinetic or pharmacodynamic co-action resulting from the combination of therapeutic agents. Administration of these therapeutic agents in combination typically is carried out over a defined time period (usually minutes, hours, days or weeks depending upon the combination selected). "Combination therapy" may, but generally is not, intended to encompass the administration of two or more of these therapeutic agents as part of separate monotherapy regimens that incidentally and arbitrarily result in the combinations contemplated by the present invention. "Combination therapy" is intended to embrace administration of these therapeutic agents in a sequential manner, that is, wherein each therapeutic agent is administered at a different time, as well as administration of these therapeutic agents, or at least two of the therapeutic agents, in a substantially simultaneous manner. Substantially simultaneous administration can be accomplished, for example, by administering to the subject a single capsule having a fixed ratio of each therapeutic agent or in multiple, single capsules for each of the therapeutic agents. Sequential or substantially simultaneous administration of each therapeutic agent can be effected by any appropriate route including, but not limited to, oral routes.

intravenous routes, intramuscular routes, and direct absorption through mucous membrane tissues. The therapeutic agents can be administered by the same route or by different routes. For example, a first therapeutic agent of the combination selected may be administered by intravenous injection while the
5 other therapeutic agents of the combination may be administered orally.

Alternatively, for example, all therapeutic agents may be administered orally or all therapeutic agents may be administered by intravenous injection. The sequence in which the therapeutic agents are administered is not narrowly critical. "Combination therapy" also can embrace the administration of the
10 therapeutic agents as described above in further combination with other biologically active ingredients and non-drug therapies (e.g., surgery or radiation treatment). Where the combination therapy further comprises a non-drug treatment, the non-drug treatment may be conducted at any suitable time so long as a beneficial effect from the co-action of the combination of the
15 therapeutic agents and non-drug treatment is achieved. For example, in appropriate cases, the beneficial effect is still achieved when the non-drug treatment is temporally removed from the administration of the therapeutic agents, perhaps by days or even weeks.

The α -aminoamide compositions of the invention can be administered in
20 a variety of dosage forms, e.g., orally, in the form of tablets, troches, capsules, sugar or film coated tablets, liquid solutions, emulsions or suspensions; rectally, in the form of suppositories; parenterally, e.g., by intramuscular or intravenous injection or infusion; and transdermally in the form of a patch, ointment, emulsion, lotion, solution, gel, cream and nasal spray.

25 Sutable pharmaceutically acceptable, therapeutically inert organic and/or inorganic carrier or excipient materials useful in the preparation of such composition include, for example, water, gelatin, gum arabic, lactose, starch, cellulose, magnesium stearate, talc, vegetable oils, cyclodextrins,

polyalkyleneglycols and the like. The α -aminoamide compositions of formula (I) can be sterilized and may contain further components, well known to those skilled in the art, such as, for example, preservatives, stabilizers, wetting or emulsifying agents, *e.g.*, paraffin oil, mannide monooleate, salts to adjust osmotic pressure, buffers and the like.

Additionally, the solid oral forms can contain, together with the active agent, diluents, *e.g.*, lactose, dextrose, saccharose, cellulose, corn starch or potato starch; lubricants, *e.g.*, silica, talc, stearic acid, magnesium or calcium stearate, and/or polyethylene glycols; binding agents, *e.g.*, starches, arabic gums, gelatin, methylcellulose, carboxymethylcellulose or polyvinyl pyrrolidone; disaggregating agents, *e.g.*, a starch, alginic acid, alginates or sodium starch glycolate; effervescing mixtures; dyestuffs; sweeteners; wetting agents such as lecithin, polysorbates, laurylsulphates; and, in general, non-toxic and pharmacologically inactive substances used in pharmaceutical formulations. The pharmaceutical preparations may be manufactured in any known manner, for example, by means of mixing, granulating, tableting, sugar-coating, or film-coating processes.

The oral formulations comprise sustained release formulations which can be prepared in a conventional manner, for instance by applying an enteric coating to tablets and granules.

The liquid dispersion for oral administration may be *e.g.*, syrups, emulsions and suspension. The syrups may further contain as a carrier, for example, saccharose or saccharose with glycerine and/or mannitol and/or sorbitol.

Suspensions and emulsions may contain as a carrier, for example, a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl alcohol. The suspensions or solutions for intramuscular injections may contain, together with the active compound, a pharmaceutically acceptable carrier, *e.g.*, sterile water, olive oil, fatty acids, glycols, *e.g.*,

propylene glycol, and, if desired, a suitable amount of lidocaine hydrochloride. The solutions for intravenous injections or infusion may contain as a carrier, for example, sterile water or preferably they may be in the form of sterile, aqueous, or isotonic saline solutions.

5 The suppositories may contain, together with the active agent, a pharmaceutically acceptable carrier, e.g., cocoa butter, polyethylene glycol, a polyoxyethylene sorbitan fatty acid ester surfactant or lecithin.

Compositions including α -aminoamides of formula (I) are generally in the form of a dose unit containing, for example, 20 to 7000 mg of active
10 ingredient per unit dosage form. Suitable treatment is given 1 or 2 or 3 times daily, depending upon clearance rate. Accordingly, the desired dose may be presented in a single dose or as divided doses administered at appropriate intervals, for example, two to four or more sub-doses per day.

The pharmaceutical compositions including an α -aminoamide of
15 formula (I) can contain, per dosage unit, e.g., capsule, tablet, powder injection, teaspoonful, suppository and the like, from about 20 to 7000 mg of the active agent.

Optimal therapeutically effective doses to be administered may be readily determined by those skilled in the art and will vary, basically, with the
20 strength of the preparation, with the mode of administration and with the advancement of the lower urinary tract disorders treated. In addition, factors associated with the particular subject being treated, including subject age, weight, diet and time of administration, will result in the need to adjust the dose to an appropriate therapeutically effective level.

25 The advantages derived from the uses and the methods of the invention as above defined are many, and include the possibility to prevent and treat basically all types of lower urinary tract disorders.

The effect of a representative compound of formula I, (S)-(+)-2-[4-(2-

fluorobenzyloxy)-benzylamino]-propanamide (NW-1029), on voiding function was tested in the following rat models of "Acute bladder irritation by acetic acid" and "Intermediate bladder irritation by cyclophosphamide (CYP)".

Continuous infusion cystometrograms (0.04 ml/min) were used to measure intercontraction intervals (ICIs) which provide an estimate of bladder capacity. Cumulative dose response curves were obtained in each animal by administering intravenously increasing doses of the drug.

Acute bladder irritation by acetic acid in rats

Procedure:

Experiments were performed using adult anesthetized female Sprague Dawley rats (170-200 g).

A catheter (PE-50) was inserted via a midline abdominal incision into the bladder through the bladder dome, and then intravesical pressure was measured to monitor bladder activity during continuous infusion of 0.15% of acetic acid. Intercontraction intervals (ICIs), maximal contraction pressure, and pressure thresholds inducing reflex bladder contraction were measured before and after intravesical infusion of acetic acid in NW-1029 treated rats.

Results:

Continuous intravesical infusion of acetic acid irritates the bladder and reduces the ICI in anesthetized rats. NW-1029 (3, 10 and 30 mg/kg) produces a significant reversal of the AA-induced reduction in the ICI in anesthetized rats (Fig. 1).

Intermediate bladder irritation by cyclophosphamide (CYP) in rats

Procedure:

Experiments were performed using both adult awake and anesthetized female Sprague Dawley rats (170-200 g).

Chemical cystitis was induced by CYP, which is metabolized to acrolein, an irritant eliminated in the urine. CYP (150 mg/kg) was

administred one day before the experiment.

NW-1029 was administered in a dose range from 0.1 to 10 mg/kg/iv.

Results:

5 Pretreatment with CYP causes bladder irritation and very frequent voidings with an ICI of about 150-200 sec. between voids. The administration of NW-1029 in a dose range from 0.1 to 10 mg/kg/iv significantly increases the ICI in both awake and anesthetized rats (Fig. 2-3).

The following EXAMPLES are presented in order to more fully illustrate the preferred embodiments of the invention.

10

EXAMPLE 1

2-[(3-Phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-propanamide

To a solution of N-methyl-alaninamide hydrochloride (0.50 g, 3.61 mmol) in methanol (10 ml), in the presence of molecular sieves (1.00 g),
15 sodium cyanoborohydride (0.36 g, 5.69 mmol) and a solution of 3-(2-phenylethyl)-2,3-dihydro-1-benzofuran-5-carboxaldehyde (0.90 g, 3.61 mmol) in methanol (10 ml) were added at room temperature. The reaction mixture was kept under stirring and an argon atmosphere for 12 h. Then, the solvent was evaporated under vacuum and the residue was purified by flash
20 chromatography on silica gel (ethyl acetate/methanol = 98/2) affording 0.93 g of title compound (77% yield).

¹H-NMR (200 MHz, CDCl₃); δ: 1.30 (d, 3H, J = 6.8 Hz); 1.76-2.19 (m, 2H); 2.64-2.87 (complex, 6H, J = 7.7 Hz); 3.29 (q, 1H, J = 7.0 Hz); 3.35-3.50 (m, 1H); 3.66 (s, 2H); 4.19-4.26 (m, 1H); 4.57-4.66 (dd, 1H, J = 9.2
25 Hz); 6.72 (d, 1H, J = 8.2 Hz); 7.01-7.32 (complex, 7H).

¹³C-NMR (200 MHz, CDCl₃); δ: 19.5; 25.9; 33.5; 36.6; 41.4; 52.3; 57.6; 77.0; 109.4; 124.4; 126.1; 128.4; 128.5; 130.8; 131.3; 141.4; 159.5; 175.0.

Analogously the following compounds were also synthesized:

2-[(3-Phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-
propanamide;

2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-propanamide;

5 2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-N-methyl-propanamide;

2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-propanamide;

10 2-{3-[2-(3-Chloro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-propanamide;

2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-
amino)-N-methyl-propanamide;

2-[(3-Phenethyl-2,3-dihydro-benzopyran-6-ylmethyl)-amino]-
propanamide;

15 2-[(4-Phenethyl-2,3-dihydro-benzoxepin-7-ylmethyl)-amino]-
propanamide;

2-[(3-Phenethyl-2,3-dihydro-benzothiophen-5-ylmethyl)-amino]-
propanamide;

20 2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide;

2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide;

2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-N-methyl-propanamide;

25 EXAMPLE 2

2-[(3-Benzyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-
propanamide

To a solution of (S)-N-methyl-glucosamine hydrochloride (0.50 g, 3.6

mmol) in methanol (10 ml), in the presence of molecular sieves (1.00 g), sodium cyanoborohydride (0.44 g, 6.9 mmol) and a solution of 3-benzyl-2,3-dihydro-1-benzofuran-5-carbaldehyde (0.75 g, 3.6 mmol) in methanol (10 ml) were added at room temperature. The reaction mixture was kept under stirring and an argon atmosphere for 12 h. Then, the solvent was evaporated under vacuum and the residue was purified by flash chromatography on silica gel (ethyl acetate) affording 0.58 g of title compound (50% yield).

¹H-NMR (400 MHz, CDCl₃); δ: 1.32 (d, 3H, J = 6.9 Hz); 2.83-2.91 (m, 4H); 3.05 (dd, J = 2.4, 6.5 Hz, 1H); 3.26 (q, 1H, J = 7.0 Hz); 3.62 (s, 2H); 3.73-3.76 (m, 1H); 4.32 (dd, 1H, J = 5.7, 8.9 Hz); 4.55 (t, 1H, J = 9.0 Hz); 6.76 (d, 1H, J = 8.1 Hz); 6.86 (bs, 1H); 7.07 (d, 1H, J = 8.1 Hz); 7.27-7.36 (m, 5H).

¹³C-NMR (400 MHz, CDCl₃); δ: 19.72; 25.84; 41.04; 43.39; 52.29; 57.55; 109.42; 124.49; 126.51; 128.35; 128.53; 129.05; 130.68; 131.16; 139.06; 159.36; 175.37.

2-[(3-Benzyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-propanamide;
2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-propanamide;

2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-N-methyl-propanamide;

2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-propanamide;

2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-N-methyl-propanamide;

2-[(3-Benzyl-2,3-dihydro-benzothiophen-5-ylmethyl)-amino]-propanamide;

2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-propanamide;

2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide.

EXAMPLES 3-6: PREPARATION OF INTERMEDIATES

EXAMPLE 3

5 3-(2-Phenylethyl)-2,3-dihydro-1-benzofuran-5-carbaldehyde

To a suspension of PCC (3.69 g, 17.10 mmol) in methylene chloride (20 ml), a solution of 3-(2-phenylethyl)-2,3-dihydro-1-benzofuran-5-yl-methanol (3.32 g, 13.16 mmol) in methylene chloride (100 ml) was added at room temperature. After stirring for 12 h, diethyl ether (300 ml) was added and the
10 black solid precipitated was filtered and washed with diethyl ether several times. The filtrate was concentrated under vacuum and the residue was purified by flash chromatography on silica gel (petroleum ether/ethyl acetate = 95/5-9/1) affording 0.202 g of the title compound (67% yield).

¹H-NMR (200 MHz, CDCl₃); δ: 1.85-2.19 (m, 2H); 2.70 (dd, 2H, J =
15 7.8 Hz); 3.44-3.56 (m, 1H); 4.33 (dd, 1H, J = 6.6, 9.2 Hz); 4.72 (dd, 1H, J = 9.2 Hz); 6.86 (d, 1H, J = 8.2 Hz); 7.10-7.34 (m, 5H); 7.63-7.69 (m, 2H); 7.72-7.73 (m, 1H); 9.82 (s, 1H).

Analogously, starting from 3-benzyl-2,3-dihydro-1-benzofuran-5-yl-methanol the 3-benzyl-2,3-dihydro-1-benzofuran-5-carboxaldehyde was
20 obtained in 85% yield.

¹H-NMR (200 MHz, CDCl₃); δ: 1.17 (t, 3H, J = 7.1 Hz); 2.77 (dd, 1H, J = 13.8, 9.4 Hz); 3.01 (dd, 1H, J = 13.9, 5.9 Hz); 3.68-3.75 (m, 1H); 4.32 (dd, J = 9.1, 5.9 Hz, 1H); 4.56 (t, J = 9.1 Hz, 2H); 6.78 (d, 1H, J = 8.3 Hz); 7.07-7.24 (m, 5H); 7.45 (bs, 1H); 7.61 (dd, 1H, J = 8.1, 1.5 Hz); 9.71 (s, 1H).

25

EXAMPLE 4

3-(2-Phenylethyl)-2,3-dihydro-1-benzofuran-5-yl-methanol

To a solution of ethyl 3-(2-phenylethyl)-2,3-dihydro-1-benzofuran-5-carboxylate (4.75 g, 14.15 mmol) in toluene (150 ml) a 1.5 M solution of

DIBAL in toluene (24 ml, 35.48 mmol) was added at -30°C. The reaction mixture was kept under stirring and an argon atmosphere at the same temperature for 30 minutes. Then, 3N HCl was added (100 ml) and the organic layer was separated. The aqueous layer was extracted with ethyl acetate (3x50 ml) and the collected organic layers were washed with saturated aqueous sodium chloride and dried over sodium sulfate. The solvent was eliminated with a rotary evaporator. The crude product (4.05 g, 99% yield) was used preceeding step without any further purification.

¹H-NMR (200 MHz, CDCl₃); δ: 1.66 (bs, 1H); 1.75-2.21 (m, 2H); 2.70 (dd, 2H, J = 7.6 Hz); 3.37-3.51 (m, 1H); 4.23 (dd, 1H, J = 6.6, 9.0 Hz); 4.58 (s, 2H); 4.63 (dd, 1H, J = 8.8 Hz); 6.74 (d, 1H, J = 8.2 Hz); 7.07-7.33 (m, 7H).

¹³C-NMR (200 MHz, CDCl₃); δ: 33.4; 36.5; 41.4; 65.4; 77.0; 109.4; 123.7; 126.1; 127.7; 128.3; 128.5; 131.2; 131.1; 141.4; 159.7.

Analogously, starting from ethyl 3-benzyl-2,3-dihydro-1-benzofuran-5-carboxylate the 3-benzyl-2,3-dihydro-1-benzofuran-5-yl-methanol was obtained in 62% yield.

¹H-NMR (200 MHz, CDCl₃); δ: 1.18 (t, 3H, J = 7.2 Hz); 2.74 (dd, 1H, J = 13.8, 9.3 Hz); 3.00 (dd, 1H, J = 13.8, 6.1 Hz); 3.51-3.70 (m, 1H); 4.21 (dd, 1H, J = 8.9, 5.9 Hz); 4.50 (m, 3H); 6.68 (d, 1H, J = 8.1 Hz); 6.93 (d, J = 0.5 Hz, 1H); 7.01-7.24 (m, 6H).

EXAMPLE 5

Ethyl 3-(2-phenylethyl)-2,3-dihydro-1-benzofuran-5-carboxylate

To a solution of ethyl 3-iodo-4-[(2E)-4-phenyl-but-2-enyl]-oxy}benzoate (13.42 g, 31.73 mmol) and AIBN (0.52 g, 3.17 mmol) in benzene (900 ml) tributyltin hydride (12 ml, 44.43 mmol) was added at room temperature and the reaction mixture was warmed to reflux. After 2 h, the solvent was eliminated under vacuum. The residue was purified by flash chromatography on silica gel (petroleum ether/ethyl acetate = 95/5) affording

9.14 g of the title compound (97% yield).

¹H-NMR (200 MHz, CDCl₃); δ: 1.36 (t, 3H, J = 7.5 Hz); 1.80-2.26 (m, 2H); 2.67 (dd, 2H, J = 7.8 Hz); 3.39-3.54 (m, 1H); 4.27-4.34 (m, 3H); 4.69 (dd, 1H, J = 9.0 Hz); 6.77 (d, 1H, J = 7.4 Hz); 7.16-7.32 (m, 5H); 7.85-7.90 (m, 2H).

¹³C-NMR (200 MHz, CDCl₃); δ: 14.4; 33.3; 36.6; 40.8; 60.6; 77.7; 109.2; 123.7; 126.1; 128.4; 128.5; 131.2; 141.2; 163.1; 166.5.

Analogously, starting from ethyl 3-iodo-4-[(2E)-3-phenyl-prop-2-enyl]-oxy]-benzoate the ethyl 3-benzyl-2,3-dihydro-1-benzofuran-5-carboxylate was obtained in 90% yield.

¹H-NMR (200 MHz, CDCl₃); δ: 1.27 (t, 3H, J = 7.1 Hz); 2.74 (dd, 1H, J = 13.8, 9.4 Hz); 3.03 (dd, 1H, J = 13.9, 5.9 Hz); 3.65-3.73 (m, 3H); 4.18-4.32 (m, 1H); 4.50 (t, J = 9.1 Hz, 2H); 6.70 (d, 1H, J = 8.3 Hz); 7.07-7.24 (m, 5H); 7.63 (bs, 1H), 7.81 (dd, 1H, J = 8.1, 1.5 Hz).

EXAMPLE 6

Ethyl 3-iodo-4-[(2E)-4-phenyl-but-2-enyl]-oxy}-benzoate

To a suspension of NaH (0.82 g, 20.5 mmol) in THF (40 ml), a solution of ethyl 4-hydroxy-3-iodo-benzoate (5.0 g, 17.1 mmol) in THF (40 ml), then 15-crown-5 (4.0 ml, 25.6 mmol) and finally a solution of 4-phenylbut-2-enyl bromide (5.41 g, 25.6 mmol) in THF (50 ml) were added subsequently at 0°C. The reaction mixture was kept under stirring at room temperature for 10 minutes and then warmed to 50°C for 6 h. After equilibration at room temperature, 3N HCl was added (30 ml) and the mixture poured into a separatory funnel. The organic layer was separated and the aqueous layer extracted with ethyl acetate (3x30 ml). The collected organic layers were washed with saturated aqueous sodium chloride and dried over sodium sulfate. The solvent was eliminated with a rotary evaporator. The residue was purified by flash chromatography on silica gel (petroleum ether/ethyl acetate = 25/5).

affording 5.94 g of the title compound (82% yield).

¹H-NMR (200 MHz, CDCl₃); δ: 1.36 (t, 3H, J = 7.2 Hz); 3.43 (d, 2H, J = 6.6 Hz); 4.33 (q, 2H, J = 7.2 Hz); 4.61 (dd, 2H, J = 1.2, 5.6 Hz); 5.67-6.15 (m, 2H); 6.78 (d, 1H, J = 8.6 Hz); 7.15-7.30 (m, 5H); 7.96 (dd, 1H, J = 2.0, 8.6 Hz); 8.44 (d, 1H, J = 2.0 Hz).


¹³C-NMR (200 MHz, CDCl₃); δ: 14.4; 38.7; 61.0; 69.6; 111.3; 125.1; 126.3; 128.5; 131.4; 134.1; 141.0.

Analogously, starting from ethyl cinnamyl bromide the ethyl 3-Iodo-4-[[*(2E)*-3-phenylprop-2-enyl]-oxy]-benzoate was obtained in 72% yield.

¹H-NMR (200 MHz, CDCl₃); δ: 1.28 (t, 3H, J = 7.1 Hz); 4.24 (q, 2H, J = 7.1 Hz); 4.69 (dd, 2H, J = 1.5-5.4 Hz); 6.24-6.35 (m, 1H), 6.68-6.76 (m, 2H); 7.16-7.34 (m, 5H); 7.90 (dd, 1H, J = 2.0, 8.6 Hz); 8.38 (d, 1H, J = 1.8 Hz).

CLAIMS

1. The use of at least one therapeutic agent which is an alpha-aminoamide compound of formula (I),



(I)

wherein:

- R is a furyl, thienyl, or pyridyl ring or a phenyl ring, optionally substituted by one or two substituents independently selected from halogen, hydroxy, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy or trifluoromethyl;
- R₁ is hydrogen or C₁-C₆ alkyl or C₃-C₇ cycloalkyl;
- R₂ and R₃ are independently selected from hydrogen; C₁-C₄ alkyl, optionally substituted by hydroxy or phenyl, phenyl, the phenyl rings being optionally substituted by one or two substituents independently selected from C₁-C₆ alkyl, halogen, hydroxy, C₁-C₆ alkoxy or trifluoromethyl; or R₂ and R₃, taken with the carbon atom which they are linked to, form a C₃-C₆ cycloalkyl ring;
- R₄, R₅ are, independently, hydrogen, C₁-C₆ alkyl or C₃-C₇ cycloalkyl; or R₄ and R₅, taken together with the nitrogen atom they are linked to, form a 5-7 atom saturated heterocyclic ring;
- X is CH₂, O, S;
- Y, Z are hydrogen or taken together form a 5-7 saturated or unsaturated carbocycle or a heterocycle;

or isomers, mixtures, and pharmaceutically acceptable salts thereof for the

preparation of a medicament for treating lower urinary tract disorders.

2. The use according to claim 1, wherein the compound is selected from:

2-[(3-Benzyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-propanamide;

2-[(3-Benzyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-

5 propanamide;

2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;

2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-N-
methyl-propanamide;

10 2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;

2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-N-
methyl-propanamide;

2-[(3-Phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-propanamide;

15 2-[(3-Phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-
propanamide;

2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;

2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-

20 N-methyl-propanamide;

2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;

2-{3-[2-(3-Chloro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;

25 2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
N-methyl-propanamide;

2-[(3-Phenethyl-2,3-dihydro-benzopyran-6-ylmethyl)-amino]-propanamide;

2-[(4-Phenethyl-2,3-dihydro-benzoxepin-7-ylmethyl)-amino]-propanamide;


- 2-[(3-Benzyl-2,3-dihydro-benzothiophen-5-ylmethyl)-amino]-propanamide;
2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-
propanamide;
2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-
5 propanamide;
2-[(3-Phenethyl-2,3-dihydro-benzothiophen-5-ylmethyl)-amino]-propanamide;
2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide;
2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
10 amino)-propanamide;
2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-N-methyl-propanamide.
3. The use according to claim 1, wherein the compound is selected from:
2-(4-Phenethyl-benzylamino)-propanamide;
15 2-(4-Benzyloxybenzylamino)-propanamide;
2-(3-Benzyloxybenzylamino)-propanamide;
2-(4-Benzylthiobenzylamino)-propanamide;
2-(4-Benzyloxy-benzylamino)-3-N-dimethyl-butanamide;
2-[4-(2-Methoxybenzyloxy)-benzylamino]-propanamide;
20 2-[4-(2-Fluorobenzyloxy)-benzylamino]-propanamide;
2-[3-(2-Fluorobenzyloxy)-benzylamino]-propanamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-propanamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-2-methyl-propanamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-N-methyl-propanamide;
25 2-[3-(3-Fluorobenzyloxy)-benzylamino]-propanamide;
2-[4-(3-Fluorobenzyloxy)-benzylamino]-propanamide;
2-[4-(3-Methoxybenzyloxy)-benzylamino]-propanamide;
2-[4-(3-Cyano-benzyloxy)-benzylamino]-propanamide;

- 2-[4-(3-Fluorobenzyloxy)-benzylamino]-propanamide;
2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-methyl-propanamide;
2-[4-(3-Fluorobenzyloxy)-benzylamino]-N-methyl-propanamide;
2-[4-(4-Fluorobenzyloxy)-benzylamino]-propanamide;
5 2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-methyl-propanamide;
2-[4-(2-Chlorobenzyloxy)-benzylamino]-propanamide;
2-[4-(3-Chlorobenzyloxy)-benzylamino]-propanamide;
2-(4-Benzyloxybenzylamino)-3-hydroxy-propanamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-3-hydroxy-propanamide;
10 2-[4-(3-Fluorobenzyloxy)-benzylamino]-3-hydroxy-propanamide;
2-(4-Benzyloxybenzylamino)-3-hydroxy-N-methyl-propanamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-3-hydroxy-N-methyl-propanamide;
2-[4-(3-Fluorobenzyloxy)-benzylamino]-3-hydroxy-N-methyl-propanamide;
2-[4-(2-Chlorobenzyloxy)-benzylamino]-3-hydroxy-N-methyl-propanamide;
15 2-[4-(3-Cyanobenzyloxy)-benzylamino]-3-hydroxy-N-methyl-propanamide;
2-[4-(3-Cyanobenzyloxy)-benzylamino]-2-methyl-3-hydroxy-N-methyl-propanamide;
2-[4-(3-Chlorobenzyloxy)-phenylethylamino]-propanamide;
2-[4-[2-(3-Fluorophenyl)-ethyloxy]benzylamino]-propanamide;
20 2-[4-[2-(3-Fluorophenyl)-ethyl]benzylamino]-propanamide;
2-[N-(4-Benzyloxybenzyl)-N-methylamino]-propanamide;
2-[4-[(3-Chlorobenzyloxy)-phenylethyl]-amino]-propanamide;
2-[4-Benzylthiobenzylamino]-propanamide;
2-[4-(2-Fluorobenzylthio)-benzylamino]-propanamide;
25 2-[4-(3-Fluorobenzylthio)-benzylamino]-propanamide;
2-[4-(3-Phenylpropyloxy)-benzylamino]-propanamide;
2-[4-(4-Phenylbutyloxy)-benzylamino]-propanamide;
2-[4-(5-Phenylpentyloxy)-benzylamino]-propanamide;

- 2-(4-Benzyloxybenzylamino)-3-phenyl-N-methyl-propanamide;
2-(4-Benzyloxybenzylamino)-3-methyl-N-methyl-butanamide;
2-(4-Benzyloxybenzylamino)-2-phenyl-acetamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-2-phenyl-acetamide;
5 2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-phenyl-acetamide;
2-[4-(2-Fluorobenzyloxy)-benzyl-N-methylamino]-2-phenyl-acetamide;
2-[4-(3-Fluorobenzyloxy)-benzyl-N-methylamino]-2-phenyl-acetamide;
2-[4-(3-Chlorobenzyloxy)-benzylamino]-2-phenyl-acetamide;
2-[4-(2-Fluorobenzyloxy)-benzylamino]-2-(2-fluorophenyl)-acetamide;
10 2-[4-(2-Fluorobenzyloxy)-benzylamino]-2-(3-fluorophenyl)-acetamide;
2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-(2-fluorophenyl)-acetamide;
2-[4-(3-Fluorobenzyloxy)-benzylamino]-2-(3-fluorophenyl)-acetamide;
2-[4-(3-Chlorobenzyloxy)-benzylamino]-2-(3-fluorophenyl)-acetamide;
2-(4-(2-Thienyloxy)-benzylamino)-propanamide.
- 15 4. The use of claim 1, wherein said lower urinary tract disorders are overactive bladder (OAB), prostatitis and prostatic dysplasia, interstitial cystitis, benign prostatic hyperplasia and urinary incontinence as the consequence of the above pathologies.
5. The use of any one of claims 1 to 4, wherein the α -aminoamides are (S)-
20 (+)-2-[3-(2-fluorobenzyloxy)-benzylamino]-propanamide; (S)-(+)-2-[4-(2-fluorobenzyloxy)-benzylamino]-propanamide, (S)-(+)-2-[4-(2-fluorobenzyloxy)-benzylamino]-N-methyl-propanamide, (S)-(+)-2-[3-(3-fluorobenzyloxy)-benzylamino]-propanamide, (S)-(+)-2-[4-(3-fluorobenzyloxy)-benzylamino]-propanamide, (S)-(+)-2-[4-(3-fluorobenzyloxy)-benzylamino]-N-methyl-propanamide, (S)-(+)-2-[4-(3-fluorobenzyloxy)-benzylamino]-N-methyl-propanamide, (S)-(+)-2-(4-phenethyl-benzylamino)-propanamide, (R)-(-)-2-(4-benzyloxy-benzylamino)-3-N-dimethyl-butanamide, (S)-(+)-2-(4-benzylthiobenzylamino)-propanamide and 2-[(S)-phenethyl-3,5-dihydro-1,4-benzodioxin-5-ylmethyl]-amino-1-N-methyl-propanamide and (S)-(+)-2-(4-

phenethyl-2,3-dihydro-benzo furan-5-ylmethyl)-amino]-N-methyl-
propanamide, 2-{3-[2-(2-fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-
ylmethyl)-amino)-propanamide, 2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-
benzothiophen-5-ylmethyl)-amino)-propanamide, (2R/3'S,R)-2-{3-[2-(2-
5 fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl)-amino)-
propanamide, (2R/3'S,R)-2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-
benzothiophen-5-ylmethyl)-amino)-propanamide.

6. An α -aminoamide compound of formula (I),



(I)

wherein:

- R is a furyl, thienyl, or pyridyl ring or a phenyl ring, optionally substituted by one or two substituents independently selected from halogen, hydroxy, cyano, C₁-C₆ alkyl, C₁-C₆ alkoxy or trifluoromethyl;
- R₁ is hydrogen or C₁-C₆ alkyl or C₃-C₇ cycloalkyl;
- R₂ and R₃ are independently selected from hydrogen; C₁-C₄ alkyl, optionally substituted by hydroxy or phenyl, phenyl, the phenyl rings being optionally substituted by one or two substituents independently selected from C₁-C₆ alkyl, halogen, hydroxy, C₁-C₆ alkoxy or trifluoromethyl; or R₂ and R₃, taken with the carbon atom which they are linked to, form a C₃-C₆ cycloalkyl ring;
- R₄, R₅ are, independently, hydrogen, C₁-C₆ alkyl or C₃-C₇ cycloalkyl; or R₄ and R₅, taken together with the nitrogen atom they are linked to, form a 5-7 atom saturated heterocyclic ring;

- X is CH₂, O, S;
- Y, Z taken together form a dihydrobenzofuran or dihydrobenzothiophen or a dihydrobenzo(thio)pyran or a tetrahydrobenz(thio)oxepin heterocycle, isomers, mixtures, and pharmaceutically acceptable salts thereof.

5 7. A compound according to claim 6, selected from:

- 2-[(3-Benzyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-propanamide;
2-[(3-Benzyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-
propanamide;
10 2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;
2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-N-
methyl-propanamide;
2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
15 propanamide;
2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-N-
methyl-propanamide;
2-[(3-Phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-propanamide;
2-[(3-Phenethyl-2,3-dihydro-benzofuran-5-ylmethyl)-amino]-N-methyl-
20 propanamide;
2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;
2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
N-methyl-propanamide;
25 2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;
2-{3-[2-(3-Chloro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
propanamide;

- 2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzofuran-5-ylmethyl}-amino)-
N-methyl-propanamide;
- 2-[(3-Phenethyl-2,3-dihydro-benzopyran-6-ylmethyl)-amino]-propanamide;
- 2-[(4-Phenethyl-2,3-dihydro-benzoxepin-7-ylmethyl)-amino]-propanamide;
- 5 2-[(3-Benzyl-2,3-dihydro-benzothiophen-5-ylmethyl)-amino]-propanamide;
- 2-{3-[2-(2-Fluoro-benzyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-
propanamide;
- 2-{3-[2-(3-Fluoro-benzyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-amino)-
propanamide;
- 10 2-[(3-Phenethyl-2,3-dihydro-benzothiophen-5-ylmethyl)-amino]-propanamide;
- 2-{3-[2-(2-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide;
- 2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-propanamide;
- 15 2-{3-[2-(3-Fluoro-phenethyl)]-2,3-dihydro-benzothiophen-5-ylmethyl}-
amino)-N-methyl-propanamide
- or isomers, mixtures, and pharmaceutically acceptable salts thereof.
8. A pharmaceutical composition comprising a pharmaceutically
acceptable excipient and, as an active agent, an effective amount of a
- 20 compound of claims 6 or 7.

ABSTRACT**ALFA-AMINOAMIDE DERIVATIVES USEFUL IN THE TREATMENT
OF LOWER URINARY TRACT DISORDERS**

5

Methods of using certain α -aminoamide derivatives to treat lower urinary tract disorders. The therapeutic agents of the invention are able to reduce or even stop the lower urinary tract disorders substantially without side effects.

10

Sheet 1/3




Fig. 1 Effect of NW-1029 in AA irritated anesthetized rats
(n=4, * $p < 0.05$ ** $p < 0.01$)

Sheet 2/3




Fig. 2 Effect of NW-1029 on CYP-irritated awake rats
(n=5, * p<0.05)

Sheet 3/3




Fig. 3 Effect of NW-1029 on CYP-irritated anesthetized rats
(n=10, * p<0.05 ** p<0.01 *** p<0.001)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☒ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☒ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.